

CMB lensing reconstruction from the WMAP 7-year data

C. Sofia Carvalho^{1,2,*}, Ismael Tereno¹, and Spyros Basilakos²

¹ *Centro de Astronomia e Astrofísica da Universidade de Lisboa,
Tapada da Ajuda, 1349-018 Lisboa, Portugal and*

² *Research Center for Astronomy and Applied Mathematics,
Academy of Athens, Soranou Efessiou 4, GR 11 527 Athens, Greece*

(Dated: January 1, 2013)

Abstract

We attempt to make a direct measurement of the weak lensing signal from the WMAP 7-year data. We apply the real-space implementation of the optimal quadratic estimator on the maps produced by the W-band Differencing Assemblies. We obtain a weak lensing amplitude parameter of $A_L = 0.98 \pm 2.50$ after correcting for several sources of bias. The significance of the estimation improves to $A_L = 1.55 \pm 0.93$ when information from WMAP simulations is used. Despite the demonstrated insensitivity of the real-space estimator to uncorrelated noise, we conclude that this detection is not statistically significant. We expect that a full-sky, higher-sensitivity experiment such as Planck will allow us to make a more significant measurement.

*Electronic address: carvalho.c@gmail.com

I. INTRODUCTION

Gravitational lensing of the cosmic microwave background (CMB) radiation is a source of information on the interaction between the CMB photons and the fluctuations in the matter density distributed along their trajectory. This interaction causes the deflection of the CMB photons by the gradient of the lensing potential ψ , which results in the creation of correlations among initially uncorrelated modes. Since the matter fluctuations are in the linear regime on large scales, the deflections are small and consequently the total deflection can be described as a sum of multiple deflections along the unperturbed trajectory (Born approximation). The correlations appear in the form of non-diagonal couplings between gradients of the temperature and gradients of the lensing potential, hence as a non-linear signal which becomes important at small scales. By measuring these correlations, we can estimate the lensing deflection $\boldsymbol{\alpha} = \nabla\psi$ and consequently the lensing convergence $\kappa = -\nabla^2\psi/2$ which is a direct measure of the projected matter density.

Detections of a lensing signal in the CMB are usually summarized in the lensing amplitude parameter A_L which describes the amplitude of the detected lensing power spectrum relative to the theoretical prediction. Thus, $A_L = 0$ indicates no lensing, while $A_L = 1$ indicates the standard Λ CDM model and General Relativity lensing theory. Recently there were reported direct detections of the lensing signal from experiments designed to look for non-linear signals in the CMB, yielding $A_L = 1.16 \pm 0.29$ from the Atacama Cosmology Telescope (ACT) data [1], and $A_L = 0.90 \pm 0.19$ from the South Pole Telescope (SPT) data [2]. Although the Wilkinson Microwave Anisotropy Probe (WMAP) was not designed to look for non-linear signals, a detection from the WMAP full-sky data would provide a full-sky measurement of the lensing signal and complement the other detections.

We anticipate some difficulties in this detection. Since the experimental noise in the WMAP data overwhelms the lensing signal, a statistically significant measurement should not be possible and indeed the first detections were made indirectly by correlating the results with galaxy counts [3, 4]. There is however a reported direct detection of the lensing signal from the WMAP data using a kurtosis estimator, finding $A_L = \{0.96 \pm 0.60, 1.06 \pm 0.69, 0.97 \pm 0.47\}$ from the data of the V, W and combined V+W bands respectively [5], whereas a more recent less-significant direct detection from WMAP data, using the optimal quadratic estimator [6], found $A_L = 1.27 \pm 0.98$ [7].

Here we base our measurement on an estimator with demonstrated insensitivity to uncorrelated noise [8] and good handling of excision of points [9], which makes it a reliable tool for the extraction of the lensing signal from the WMAP data. Similarly to the optimal estimator, this estimator combines two filters of the temperature map weighted by the inverse of the variance. Instead of acting in Fourier space and using all the available sky for the reconstruction of the lensing potential at any point, this estimator acts in real space and uses the neighbouring points only, which presupposes a truncation of the kernel that convolves the two filters. Although truncating the kernel renders the estimator slightly suboptimal, it has

a strong physical motivation. Since weak lensing is a small-scale effect coherent over scales of a few degrees [10], information on the lensing potential at a given point is contained in the points within the degree-scale, so we expect little gain in considering the information over the entire sky [11]. Thus a direct detection of a lensing signal from the WMAP data is both a test of the robustness of the real-space estimator and a comparison of the performance with other estimators.

The manuscript is organized as follows. In Sec. II we introduce the weak lensing estimator. In Sec. III we justify our choice of data and describe the simulations. We then proceed to measure the weak lensing signal. In Sec. IV we use synthesised map so as to simulate a control experiment, test the bias removal and setup the calibration. In Sec. V we use patches from cartesian projections of the full-sky maps. The results are summarized in Sec. VI.

II. ESTIMATOR OF THE CMB WEAK LENSING

The deflection of the CMB photons by the gradient of the lensing potential amounts to remapping the unlensed temperature anisotropies T to the lensed $\tilde{T}$ in the direction $\boldsymbol{\theta}$ by $\tilde{T}(\boldsymbol{\theta}) = T(\boldsymbol{\theta} + \nabla\psi)$, where the lensing potential ψ is the projection on the image plane of the gravitational potential of the large-scale structure. For small deflections, the lensed temperature power spectrum is

$$\langle \tilde{T}(\boldsymbol{\ell}') \tilde{T}(\boldsymbol{\ell} - \boldsymbol{\ell}') \rangle = (2\pi)^2 \delta(\boldsymbol{\ell}) C_{\boldsymbol{\ell}'} + (2\pi)^2 [\boldsymbol{\ell} \cdot \boldsymbol{\ell}' C_{\boldsymbol{\ell}'} + \boldsymbol{\ell} \cdot (\boldsymbol{\ell} - \boldsymbol{\ell}') C_{|\boldsymbol{\ell} - \boldsymbol{\ell}'|}] \psi(\boldsymbol{\ell}) + O(\psi^2), \quad (1)$$

where $C_{\boldsymbol{\ell}}$ is the unlensed temperature power spectrum. Both the unlensed temperature anisotropies and the large-scale structure are assumed to be Gaussian random fields. The remapping of one Gaussian field by another induces non-Gaussianities. The lensing signal is encoded in the off-diagonal correlations as a coupling between the photons and the gradient of the lensing potential. An estimator that captures the information in the coupling consists of a quadratic combination of the temperature map optimized so as to yield the minimum variance [6]. This estimator will also pick up the diagonal correlations which exist in the absence of lensing and which we must remove both from the estimated map and from the estimated power spectrum. The diagonal correlations have a much larger amplitude than the lensing signal and are estimated from simulated unlensed temperature maps.

We measure the isotropic component of the convergence tensor, defined as $\kappa = -(\partial_x^2 + \partial_y^2)\psi/2$. We use the real-space estimator introduced in Ref. [8] which is based on the convolution of the square of the lensed temperature map $\tilde{T}$ with a local kernel W as follows

$$\hat{\kappa}(\boldsymbol{\theta}) = \int d^2\boldsymbol{\theta}_+ \int d^2\boldsymbol{\theta}_- \tilde{T}(\boldsymbol{\theta}_+ + \boldsymbol{\theta}_-) \tilde{T}(\boldsymbol{\theta}_+ - \boldsymbol{\theta}_-) W(\boldsymbol{\theta}, \boldsymbol{\theta}_+, \boldsymbol{\theta}_-). \quad (2)$$

The real-space kernel $W(\boldsymbol{\theta}, \boldsymbol{\theta}_+, \boldsymbol{\theta}_-)$ is given by

$$W(\boldsymbol{\theta}, \boldsymbol{\theta}_+, \boldsymbol{\theta}_-) = \int \frac{d^2\boldsymbol{\ell}_+}{(2\pi)^2} \exp[i\boldsymbol{\ell}_+ \cdot \boldsymbol{\theta}] \int \frac{d^2\boldsymbol{\ell}_-}{(2\pi)^2} \exp[-i(\boldsymbol{\ell}_+ \cdot \boldsymbol{\theta}_+ + \boldsymbol{\ell}_- \cdot \boldsymbol{\theta}_-)] W(\boldsymbol{\ell}_+, \boldsymbol{\ell}_-) \quad (3)$$

where the corresponding Fourier-space kernel $W(\ell_+, \ell_-)$ is related to the minimum-variance (to leading order) weight function $Q(\ell, \ell')$ by

$$W(\ell_+, \ell_-) = Q(\ell, \ell') \quad (4)$$

upon the coordinate transformation $\ell = \ell_+$, $\ell' = (\ell_+ + \ell_-)/2$. (See Ref. [8] for details). Here

$$Q(\ell, \ell') = \frac{\ell^2}{4} \mathcal{N}_\ell \frac{1}{2} \frac{\ell \cdot \ell' C_{\ell'} + \ell \cdot (\ell - \ell') C_{|\ell' - \ell|}}{[\tilde{C}_{\ell'} + N_{\ell'}][\tilde{C}_{|\ell - \ell'|} + N_{|\ell - \ell'|}]} \quad (5)$$

where N_ℓ is the noise power spectrum and $\mathcal{N}_\ell$ is the variance of the estimator defined by

$$\mathcal{N}_\ell = \int \frac{d^2 \ell'}{(2\pi)^2} \frac{1}{2} \frac{[\ell \cdot \ell' C_{\ell'} + \ell \cdot (\ell - \ell') C_{|\ell' - \ell|}]^2}{[\tilde{C}_{\ell'} + N_{\ell'}][\tilde{C}_{|\ell - \ell'|} + N_{|\ell - \ell'|}]} \quad (6)$$

In the ℓ -integration we sum modes $2 \leq \ell, \ell' \leq 1200$. The validity range of the real-space estimator is defined by the largest and smallest lensing modes that can be probed by the kernel. Hence, the lower ℓ -limit is determined by the size of the maps, whereas the upper ℓ -limit is determined by the size of the kernel.

III. DATA AND SIMULATIONS

The relevant information for the reconstruction of the lensing potential lies on the angular scales close to the resolution scale of the map, which is set by the beam size. Hence we opt to use the maps produced by the four W-band differential arrays (DA) for having the smallest beam size¹. To minimize the contamination of the lensing signal by spurious sources of non-Gaussianity, we use the coadded, foreground-removed temperature maps [12] at resolution $r9^2$. Moreover, to exclude foreground-contaminated portions of the sky at the location of the galactic plane and point sources, we use the pixel mask KQ85 which keeps 78% of the sky area³.

We conceive different experiments based on the specifications of the four DAs of the WMAP W-band⁴. For each differential array, we conceive two experiments: a) the experiment denoted by “theo,” which consists of lensed CMB maps synthesised in the flat-sky approximation using the temperature (TT) and lensing deflection ($\alpha\alpha$) power spectra computed with CAMB [13] (on scales $\ell \leq 1200$, for a fiducial flat Λ CDM cosmology consisting of $h = 0.7$, $\Omega_{\text{cdm}} h^2 = 0.122$, $\Omega_b h^2 = 0.0226$, $n_s = 0.96$, $\sigma_8 = 0.85$, $\tau_{\text{rei}} = 0.09$ and $T_0 = 2.725$ K), and b) the experiment denoted by “obs,” which consists of temperature maps derived from HEALPix⁵ [14] cartesian projections of the full-sky map produced by WMAP. The specifications of the experiments are summarized in Table I.

¹ http://lambda.gsfc.nasa.gov/product/map/dr4/m_products.cfm

² http://lambda.gsfc.nasa.gov/product/map/dr4/maps_da_r9_iqr_7yr_get.cfm

³ http://lambda.gsfc.nasa.gov/product/map/dr4/ptsrc_catalog_get.cfm

⁴ http://lambda.gsfc.nasa.gov/product/map/dr4/beam_xfer_get.cfm

⁵ <http://healpix.jpl.nasa.gov/>

Experiment	θ_{fwhm} (arcmin)	σ_0 ($\mu K/N_{\text{obs}}$)	$\langle N_{\text{obs}} \rangle_{\text{pix}}$	FOV_{map} (deg)	θ_{kernel} (deg)
WMAP $_{W_1^{\text{theo/obs}}}$	13.2	5.906×10^3	2.512×10^3	56	1.1
WMAP $_{W_2^{\text{theo/obs}}}$	13.2	6.572×10^3	2.515×10^3	56	1.1
WMAP $_{W_3^{\text{theo/obs}}}$	13.2	6.941×10^3	2.514×10^3	56	1.1
WMAP $_{W_4^{\text{theo/obs}}}$	13.2	6.778×10^3	2.517×10^3	56	1.1

Table I: **The specifications of the experiments used in this study.** The experiments denoted by WMAP $_{W_i}^{\text{theo}}$ use the power spectra computed with CAMB and detector noise characteristic of the W_i DA. The experiments denoted by WMAP $_{W_i}^{\text{obs}}$ use cartesian projections of the corresponding W_i DA measured by WMAP.

IV. MEASUREMENT OF THE CONVERGENCE FROM THE CMB POWER SPECTRUM

A. The input maps

We first compute the convergence map from simulations of the four DAs of the WMAP W-band. We synthesise lensed CMB temperature maps using the theoretical power spectra (TT and $\alpha\alpha$) computed with CAMB and combining the beam window function and detector noise characteristic of each DA of the WMAP W-band. (See Appendix B of Ref. [8] for details.) We compute the noise power spectrum from a full-sky map synthesised as White Noise $\times \sigma_0/\sqrt{N_{\text{obs}}}$, where σ_0 is the rms of the noise per number of observations and N_{obs} is the corresponding map containing the number of observations per pixel. From the noise power spectrum, we synthesise a noise map which we add to the beam-convolved, lensed CMB map. We call these experiments WMAP $_{W_i}^{\text{theo}}$.

For each experiment, we also generate unlensed CMB temperature maps from the theoretical TT power spectrum computed with CAMB. The unlensed CMB maps are generated using different realizations of the CMB map for the same cosmology, thus simulating different realizations of the temperature and noise maps. The unlensed CMB maps serve to model the Gaussian contribution of the CMB to the four-point correlations measured by the estimator of the lensing power spectrum which is the major contaminant of the lensing signal [15].

B. The convergence maps

The convergence map is reconstructed by convolving in real space the minimum variance kernel $W(\boldsymbol{\theta}, \boldsymbol{\theta}_+, \boldsymbol{\theta}_-)$ with the square of the temperature map from which we want to estimate the weak lensing signal. Both the lensed CMB power spectrum and a fiducial power spectrum for the unlensed CMB power spectrum enter in the calculation of the kernel. We generate




Figure 1: **Kernel in real space for the four DA's of the WMAP W-band (theo).** We plot the dominant ($m = 0$) eigenfunction only. The levels denote the orders of magnitude as fractions of the maximum value, delimited by $\{1, 10^{-1}, 10^{-2}, 10^{-3}, 10^{-4}\}$. The kernels in all panels are derived from the theoretical power spectra combined with the beam window function of each DA. Top left: W_1 , Top right: W_2 , Bottom left: W_3 , Bottom right: W_4 .

the kernel for each DA. In the calculation of the kernel we use the lensed power spectrum computed with CAMB combined with the window beam function and the detector noise characteristic of each DA.

The kernel can be expressed as a series of eigenfunctions, the dominant being the $m = 0$ mode which we plot in Fig. 1. (See Ref. [8] for the calculation of the kernel in real space.) The plots show that the kernels for the four experiments have the same size and similar structures within the 1% level, differing only in the smaller-scale structures (of order $< 0.1^\circ$) at sub-percent level, so we expect that the corresponding lensing reconstructions have similar ranges of validity. In all cases, for the numerical implementation we truncate the kernel at $\theta_{\text{kernel}} = 1.1^\circ$, which is the spatial extent which corresponds to the 1% level. As previously tested, keeping sub 1% levels does not improve the reconstruction of the power spectrum [8].

The convergence map $\hat{\kappa}$ reconstructed from a lensed CMB map contains both the lensing signal $\kappa|_\psi$ and a spurious contribution $\kappa|_{\psi=0}$ due to the unlensed CMB map. The zero-point

function of this contribution amounts to an offset which we remove by subtracting the pixel average of the reconstructed convergence map $\hat{\kappa}$. The resulting convergence map $\hat{\kappa} - \langle \hat{\kappa} \rangle_{\text{pixel}}$ is unbiased in the sense that its expectation value is equal to that of $\kappa_{|\psi}$. (See Refs. [8, 9] for details.)

C. The convergence power spectrum

After correcting for the bias of the reconstructed convergence map, spurious Gaussian correlations due to the unlensed CMB still persist which add to the non-Gaussian correlations due to the lensing signal. The power spectrum of the convergence due to the lensing signal is obtained after subtracting the Gaussian contribution off of the power spectrum of the reconstructed convergence map. To estimate the Gaussian contribution, we take the average of the power spectrum of the convergence maps reconstructed from various maps of the unlensed CMB. The unlensed CMB maps are synthesised from the same theoretical TT power spectrum while varying the seed. We estimate the unbiased power spectrum of the convergence $C_\ell^{\kappa_{|\psi}}$ by considering the weighted average of $C_\ell^{\hat{\kappa}_{|\psi=0}}$ over the different CMB realizations

$$\langle C_\ell^{\hat{\kappa}_{|\psi=0}} \rangle_{CMB} = \sum_i^{\text{num maps}} \left(C_\ell^{\hat{\kappa}_{|\psi=0}} \right)_i w_i^{\hat{\kappa}_{|\psi=0}} / \sum_i^{\text{num maps}} w_i^{\hat{\kappa}_{|\psi=0}} \quad (7)$$

and subtracting it off of $C_\ell^{\hat{\kappa}}$. The weights $w_i^{\hat{\kappa}_{|\psi=0}}$ are the inverse of the variance of the estimated power spectrum

$$\text{Var}^{\text{est}}[C_\ell^{\hat{\kappa}_{|\psi=0}}]_i = \frac{1}{\ell f_{\text{sky}}} \left[\text{Var}[\hat{\kappa}] + \left(C_\ell^{\hat{\kappa}_{|\psi=0}} \right)_i \right]^2 \quad (8)$$

and $\text{Var}[\hat{\kappa}]$ is the variance of the estimator given by Eqn. (6). The unbiased power spectrum estimated by this method is thus given by

$$C_\ell^{\kappa_{|\psi}} = C_\ell^{\hat{\kappa}} - \langle C_\ell^{\hat{\kappa}_{|\psi=0}} \rangle_{CMB}, \quad (9)$$

which is a difference between two very similar quantities. The total variance is given by

$$\text{Var}[C_\ell^{\kappa_{|\psi}}] = \text{Var}[C_\ell^{\hat{\kappa}}] + \text{Var}[C_\ell^{\hat{\kappa}_{|\psi=0}}]. \quad (10)$$

Here

$$\text{Var}[C_\ell^{\hat{\kappa}}] = \text{Var}^{\text{est}}[C_\ell^{\hat{\kappa}}], \quad (11)$$

$$\text{Var}[C_\ell^{\hat{\kappa}_{|\psi=0}}] = \left[\sum_i^{\text{num maps}} \left(1/\text{Var}^{\text{est}}[C_\ell^{\hat{\kappa}_{|\psi=0}}]_i \right) \right]^{-1} + \text{Var}^{\text{s}}[C_\ell^{\hat{\kappa}_{|\psi=0}}], \quad (12)$$

where $\text{Var}^{\text{est}}[C_\ell^{\hat{\kappa}}]$ is given by the analogous to Eqn. (8) upon the replacement $\hat{\kappa}_{|\psi=0} \rightarrow \hat{\kappa}_{|\Psi}$, and Eqn. (12) includes the sample variance about the average value of $C_\ell^{\hat{\kappa}_{|\psi=0}}$, $\text{Var}^{\text{s}}[C_\ell^{\hat{\kappa}_{|\psi=0}}]$.

The estimate of the convergence power spectrum averaged over the four experiments $\text{WMAP}_{W_i}^{\text{theo}}$ is plotted in Fig. 2 (left panel). The sample variance becomes subdominant in the entire ℓ -range with the increase of the number of maps. The detection error is dominated by the estimator noise set by the first term in Var^{est} , which depends on the experimental resolution and detector noise. We observe, however, that the result contains a large excess of power as compared to the expected signal, the input convergence power spectrum also shown in Fig. 2. In contrast, for simulations of the higher signal-to-noise Planck experiment, the quantity $\langle C_\ell^{\hat{\kappa}|\psi=0} \rangle_{\text{CMB}}$ was found to be a good estimator of the Gaussian bias as demonstrated in Refs. [8, 9].

A source of additional bias are couplings of lensing modes which add spurious correlations to the estimated power spectrum from the four-point correlation function of the lensed CMB [15]. We expect to get a much better agreement if we are able to reduce the uncertainty in the CMB map since the couplings add incoherently [15]. This was not required for the signal-to-noise ratio of Planck, but it is needed for WMAP. We then produce various lensed CMB maps, all from the same CAMB lensing and temperature power spectra, and again generate unlensed CMB maps using the same realizations of the lensed CMB maps. Throughout the different realizations we generate different realizations of the noise map.

We estimate the unbiased power spectrum of the convergence $C_\ell^{\kappa|\psi}$ by taking the difference of the weighted averages over the different CMB realizations

$$C_\ell^{\kappa|\psi} = \langle C_\ell^{\hat{\kappa}} \rangle_{\text{CMB}} - \langle C_\ell^{\hat{\kappa}|\psi=0} \rangle_{\text{CMB}}. \quad (13)$$

Here

$$\langle C_\ell^{\hat{\kappa}} \rangle_{\text{CMB}} = \sum_i^{\text{num maps}} (C_\ell^{\hat{\kappa}})_i w_i^{\hat{\kappa}} / \sum_i^{\text{num maps}} w_i^{\hat{\kappa}} \quad (14)$$

with the weights $w_i^{\hat{\kappa}}$ given by the inverse of the variance of the power spectrum $\text{Var}[C_\ell^{\hat{\kappa}}]_i$, and $\langle C_\ell^{\hat{\kappa}|\psi=0} \rangle_{\text{CMB}}$ given by Eqn. (7). The total variance of $C_\ell^{\kappa|\psi}$ is given by Eqn. (10), where $\text{Var}[C_\ell^{\hat{\kappa}}]$ is given by

$$\text{Var}[C_\ell^{\hat{\kappa}}] = \left[\sum_i^{\text{num maps}} (1/\text{Var}^{\text{est}}[C_\ell^{\hat{\kappa}}]_i) \right]^{-1} + \text{Var}^{\text{s}}[C_\ell^{\hat{\kappa}}] \quad (15)$$

and $\text{Var}[C_\ell^{\hat{\kappa}|\psi=0}]$ is given by Eqn. (12).

When applying the estimator of Eqn. (13), we indeed obtain a convergence power spectrum, also shown in Fig. 2 (left panel), that closely reproduces the input power spectrum within the validity range of the real-space estimator.

As a measure of the statistical significance of the detection, we compute the best-fit lensing amplitude parameter A_L . We fit the estimated power spectrum with functions of the form $A_L C_\ell^{\text{in}}$, where C_ℓ^{in} is the input fiducial convergence power spectrum. The χ^2 -minimization yields an analytical expression for A_L , which is essentially an inverse-variance




Figure 2: **Estimated convergence power spectrum from CMB simulations of the WMAP’s W-band DAs.** Left panel: The gray diamonds are the convergence power spectrum computed as the difference between one lensed CMB map and the inverse-variance average over different realizations of unlensed CMB maps. The black squares are the convergence power spectrum computed as the difference between the inverse-variance average over different realizations of paired lensed CMB and unlensed CMB maps. For each DA, the maps are synthesised from the theoretical power spectra combined with the beam window function and noise power spectrum of the corresponding DA. The dotted line is the theoretical convergence power spectrum computed with CAMB. The dash-dotted line is the inverse-sum of the variance of the estimator $\text{Var}[\hat{\kappa}_{|\psi}]$ for each DA. Right panel: The black line shows the ratio between the two quantities in the left panel.

weighting of the ratio $C_\ell^{\text{out}}/C_\ell^{\text{in}}$ summed over all scales ℓ [16],

$$A_L = \frac{\sum_\ell C_\ell^{\text{in}} C_\ell^{\text{out}} / \text{Var}[C_\ell^{\text{out}}]}{\sum_\ell (C_\ell^{\text{in}})^2 / \text{Var}[C_\ell^{\text{out}}]}. \quad (16)$$

The error of A_L is computed by propagating the error of Eqn. (16), yielding

$$\text{Var}[A_L] = \frac{1}{\sum_\ell (C_\ell^{\text{in}})^2 / \text{Var}[C_\ell^{\text{out}}]}. \quad (17)$$

We compute A_L for each DA, $(A_L^{\text{theo}})_{W_i}$, for the average of the four DA’s auto-correlations, $(A_L^{\text{theo}})_{\langle W_i W_i \rangle}$, and for the full combination of DA’s considering all auto and cross-correlations, $(A_L^{\text{theo}})_{\langle W_i W_j \rangle}$.

The results for the four experiments, separately and combined, in the validity range of the estimator, are presented in Table II. For the combined result we find $A_L = 1.55 \pm 0.93$, which is similar to the findings in Ref. [7].

	$(A_L^{\text{theo}})_{W_1}$	$(A_L^{\text{theo}})_{W_2}$	$(A_L^{\text{theo}})_{W_3}$	$(A_L^{\text{theo}})_{W_4}$	$(A_L^{\text{theo}})_{\langle W_i W_i \rangle}$	$(A_L^{\text{theo}})_{\langle W_i W_j \rangle}$
$\ell \in [6, 262]$	1.49 ± 2.95	1.49 ± 3.73	1.55 ± 3.90	1.59 ± 3.45	1.53 ± 1.72	1.55 ± 0.93

Table II: A_L^{theo} from each W-band DA and from the combined W-band DAs.

V. MEASUREMENT OF THE CONVERGENCE FROM THE CMB MAPS

We now turn to the problem of detecting a lensing signal in the WMAP 7-year data. Having one full-sky realization only of the lensed map, we cannot apply the estimator defined in Eqn. (13). We can however use the simulation results to infer an empirical calibration. We define the empirical calibration as the ratio between the two $C_\ell^{\kappa|\psi}$ estimated in Eqns. (9) and (13), and plot it in Fig. 2 (right panel).

A. The input maps

We then proceed to compute the convergence map from the temperature maps produced by the four DA's of WMAP W-band. We use the foreground-reduced maps and apply the KQ85 mask which keeps 78% of the sky area. We then make cartesian projections of the full-sky masked maps into patches of the same size ($56^\circ \times 56^\circ$) as the simulated maps in the previous section. The projections cover the whole sphere in steps of 45 degrees along the latitude and along the longitude. In a similar way, we also project the mask and the map containing the number of observations per pixel for each DA which we use to generate the noise patches. We call these experiments $\text{WMAP}_{W_i}^{\text{obs}}$. In Ref. [17], a similar strategy is adopted to estimate the bispectrum from a full-sky map of a Planck simulation.

For each patch we generate unlensed CMB maps. These maps consist of the same CMB map combined with the DA's beam window function, plus a noise map computed as $\text{White Noise} \times \sigma_0 / \sqrt{N_{\text{obs}}}$, where σ_0 is the rms of the noise per number of observations and N_{obs} is the corresponding patch containing the number of observations per pixel. Finally we apply the corresponding mask patch.

B. The convergence maps

We generate the kernel for each DA. In the calculation of the kernel we use the lensed CMB power spectrum measured by WMAP, combined with the window beam function and the detector noise characteristic of the DA. For the fiducial unlensed power spectrum we use the power spectrum computed with CAMB. We plot the $m = 0$ mode in Fig. 3. All kernels have similar size and similar structures within the 1% level, differing only in the smaller scale structures (of order $< 0.1^\circ$) at sub-percent level. For the numerical implementation we truncate the kernel at the 1% level, which corresponds to $\theta_{\text{kernel}} = 1.1^\circ$ similarly to the




Figure 3: **Kernel in real space for the four DA's of the WMAP W-band (obs).** We plot the dominant ($m = 0$) eigenfunction only. The levels denote the orders of magnitude as fractions of the maximum value, delimited by $\{1, 10^{-1}, 10^{-2}, 10^{-3}, 10^{-4}\}$. The kernels in all panels are derived from the WMAP TT power spectrum combined with the beam window function of each DA. Top left: W_1 , Top right: W_2 , Bottom left: W_3 , Bottom right: W_4 .

experiments in the previous section. From each patch i of the WMAP map from the W-band DA W_j , we compute the corresponding convergence map $\hat{\kappa}_{W_j}^i$. For illustration, we show three patches of the W_1 DA map obtained from projections centered at the galactic coordinates $(\text{long}, \text{lat}) = \{(0, 45^\circ\text{N}), (0, 0), (0, 45^\circ\text{S})\}$, and the corresponding convergence patches (Fig. 4). The contours of the mask are closely reproduced in the convergence patches, which demonstrates that the reconstruction is local.

C. The convergence power spectrum

To estimate the spurious correlations due to the unlensed CMB, we compute the convergence map from various realizations of the unlensed patch corresponding to the single realization of the observed patch. The average power spectrum of these convergence maps, weighted by the inverse of their variance according to Eqn. (7), is our estimate of the Gaussian contribution of the CMB to the convergence power spectrum of the patch. Thus, for




Figure 4: **Projections of the W_1 DA map and KQ85 mask centred at $(\text{long}, \text{lat}) = \{(0, 45^\circ\text{N}), (0, 0), (0, 45^\circ\text{S})\}$ respectively first, second and third row, and the resulting convergence patches.** Left panels: The masked CMB patches. Central panels: The mask patches, with amplitude 0 on the black regions and amplitude 1 elsewhere. Right panels: The $\hat{\kappa} - \langle \hat{\kappa} \rangle_{\text{pixel}}$ patches obtained from the corresponding masked CMB patches.

each patch i and for each DA W_j , we have

$$(C_\ell^{\kappa|\psi})_{i,W_j W_{j'}} = (C_\ell^{\hat{\kappa}})_{i,W_j W_{j'}} - \left(\langle C_\ell^{\hat{\kappa}\hat{\psi}=0} \rangle_{CMB} \right)_{i,W_j W_{j'}}. \quad (18)$$

For each patch, we compute the average over the four DA's, which includes four auto-correlations and six cross-correlations in both the $C_\ell^{\hat{\kappa}}$ and the $\langle C_\ell^{\hat{\kappa}\hat{\psi}=0} \rangle_{CMB}$ terms, weighted by the inverse of the variance of each $(C_\ell^{\kappa|\psi})_{i,W_j W_{j'}}$. This operation yields $(C_\ell^{\kappa|\psi})_i$. We then compute the average over all the patches, weighted by the variance of each $(C_\ell^{\kappa|\psi})_i$, yielding the final result $C_\ell^{\kappa|\psi}$.

We recall that masking a map causes a degradation in the amplitude of the estimated convergence power spectrum in comparison to the power spectrum estimated from the complete




Figure 5: **Dispersion of the estimated convergence power spectrum as a function of the lensed CMB realization used to define the empirical calibration.** The black dashed line is the raw estimated convergence power spectrum obtained from the CMB patches after subtracting the convergence power spectrum from unlensed CMB realizations. The black filled line is the convergence power spectrum obtained after calibrating the black dashed line with the realization of Fig. 2, whereas the grey lines are the spectra obtained after calibrating with the various lensed CMB realizations.

map. This amplitude degradation has been quantified for masks of point sources with varying density and radius [9]. We correct the convergence power spectrum of each patch by the amplitude degradation inferred for the area of the corresponding mask patch, predominantly consisting of contiguous large areas.

Finally we apply the empirical calibration of Fig. 2 (right panel) derived from the simulations. We remind that the calibration allows to move from the convergence power spectrum estimated from one map to the power spectrum estimated from the mean of a distribution of maps. In the previous section, this operation introduced a small extra source of uncertainty, the term $\text{Var}^s[C_\ell^{\hat{\kappa}}]$ in Eqn. (15). This term accounts for the error on the mean of the distribution and is subdominant as mentioned earlier on. In this section, we are limited to one realization that is not necessarily the mean of a distribution of an ensemble of WMAP-like realizations. To be conservative, we apply the empirical calibration with an uncertainty containing the sample variance per element of the distribution (as opposed to the uncertainty on the mean), which is propagated into the calibrated power spectrum. This gives an important contribution to the error budget, even though the detection is still dominated by the estimator noise as in the simulated case. This contribution is illustrated in Fig. 5, which shows the dispersion of the estimated convergence power spectrum if corrected by an ensemble of empirical calibrations that could be defined by each realization.

The results, before and after the empirical calibration, are presented in Fig. 6, for both the convergence and the deflection field (which is the quantity most commonly used, for easier comparison with the results from other detections). After correcting with the empirical




Figure 6: **Estimated convergence power spectrum from the CMB maps of the WMAP’s W-band DAs.** The grey diamonds are the weighted average of the difference between the power spectrum measured from the patches and the power spectrum measured from the simulations for each DA and for each patch. The black squares are the same quantity multiplied by the empirical calibration. The dotted line is the theoretical convergence power spectrum computed with CAMB. The dash-dotted line is the inverse-sum of the variance of the estimator $\text{Var}[\hat{\kappa}_{|\psi}]$ for each DA. Left panel: The convergence power spectrum. Right panel: The deflection power spectrum.

calibration of Fig. 2 (right panel), we recover a lensing signal which agrees with the theoretical prediction. In terms of the lensing amplitude parameter, after applying the empirical calibration, including the calibration dispersion in the error budget, we find a conservative value of $A_L = 0.98 \pm 2.50$ for the case of full combination of DA’s.

D. Null test: Measurement of the convergence from noise maps

We proceed to apply the estimator to noise patches synthesised for each DA from the corresponding σ_0 and N_{obs} . These are the same maps that we have added to the simulated unlensed CMB maps. For each DA, we average the convergence power spectrum over the different patches and present the results in Fig. 7.

The results are consistent with the absence of signal, showing that the estimator is free of systematics. We notice that this result did not require a calibration, highlighting the lensing origin of the spurious correlations, which are not present in this test on noise maps.

VI. CONCLUSIONS

In this manuscript we present a direct detection of the weak lensing from the WMAP 7-year data. We used the real-space variant of the optimal quadratic estimator originally defined in Fourier space.




Figure 7: **Estimated convergence power spectrum from noise maps of the WMAP’s W-band DAs.** The symbols are the convergence power spectrum averaged over the different patches for each DA. Diamond: W_1 . Cross: W_2 . Plus: W_3 . Triangle: W_4 . The squares are the estimated convergence power spectrum from the CMB maps (after the empirical calibration), which we include for comparison. The dotted line is the theoretical convergence power spectrum computed with CAMB.

First we applied the estimator to maps synthesised so as to simulate the WMAP W-band. The conventional estimator of the Gaussian bias failed to recover the input convergence power spectrum, in contrast with its success in Planck simulations. This failure signaled the presence of a non-Gaussian bias. A variation of this estimator was conceived which successfully accounted for the additional bias and recovered the input convergence power spectrum, with a lensing parameter of $A_L = 1.55 \pm 0.93$. The detection error is dominated by the variance of the estimator which depends on the experimental resolution and detector noise. By comparing the results of the two methods, we derived an empirical calibration which estimates the additional bias as a function of ℓ .

We proceeded to apply the estimator to maps obtained from HEALPix cartesian projections of the full-sky maps produced by the WMAP W-band. We used the conventional estimator of the Gaussian bias, including the correction of amplitude degradation due to masking, and the empirical calibration function. We achieved a good agreement of the detection with the theoretical prediction, as well as with the detection from the simulations, yielding $A_L = 0.98 \pm 2.50$. This detection is also compatible with previous detections. However, the large error renders this detection not statistically significant, similarly to the detection from the WMAP data using the harmonic implementation of optimal estimator [7]. The detection from the WMAP data using a kurtosis estimator has higher significance because it aims for the lensing correlations of interest [5].

Nonetheless, our detection serves as a proof of the robustness of the real-space estimator in dealing with extended masks and overwhelming experimental noise. We expect that a

full-sky experiment with higher resolution and lower detector noise, such as Planck [18], will allow us to make a more significant measurement of the convergence power spectrum.

Acknowledgments CSC is funded by Fundação para a Ciência e a Tecnologia (FCT), Grant no. SFRH/BPD/65993/2009. IT is funded by FCT, Grant no. SFRH/BPD/65122/2009 and acknowledges support from the European Programme FP7-PEOPLE-2010-RG-268312. CSC and IT also acknowledge support from the project CERN/FP/123618/2011. The authors acknowledge the use of the HEALPix package for the handling of the full-sky maps. The authors thank C. Feng, M.G.R. Santos, D.N. Spergel and E. Vagenas for useful discussions.

-
- [1] S. Das et al, Atacama Cosmology Telescope: Detection of the Gravitational Lensing Convergence Power Spectrum Phys. Rev. Lett. 104, 021301 (2011) (arXiv:1103.2124 [astro-ph.CO])
 - [2] A. van Engelen et al, A Measurement of Gravitational Lensing of the Microwave Background using South Pole Telescope Data (arXiv:1202.0546 [astro-ph.CO])
 - [3] K. Smith, O. Zahn and O. Doré, Detection of the Gravitational Lensing in the Cosmic Microwave Background Phys. Rev. D. 76, 043510 (2007) (arXiv:0705.3980 [astro-ph])
 - [4] C. Hirata, S. Ho, N. Padmanabhan, U. Seljak and N. Bahcall, Correlation of CMB with large-scale structure: II. Weak lensing Phys. Rev. D. 78, 043520 (2008) (arXiv:0801.0644 [astro-ph])
 - [5] J. Smidt, A. Cooray, A. Amblard, S. Joudaki, D. Munshi, M.G. Santos and P. Serra, A Constraint on the Integrated Mass Power Spectrum Out to $z = 1100$ from Lensing of the Cosmic Microwave Background, Astrophys. J. Lett. 728, 1 (2011) [arXiv:1012.1600 (astro-ph)]
 - [6] W. Hu, Mapping the dark matter through the cosmic microwave background damping tail, Astrophys. J. 557, L79, (2001) (arXiv: astro-ph/0105424)
 - [7] C. Feng, B. Keating, H.P. Paar and O. Zahn, Reconstruction of the gravitational lensing using WMAP 7-year data Phys. Rev. D 85, 043513 (2012) (arXiv:1111.2371 [astro-ph.CO])
 - [8] C.S. Carvalho and K. Moodley, Real space estimator for the weak lensing convergence from the CMB, Phys. Rev. D 81, 123010 (2010) (arXiv:1005.4288 [astro-ph])
 - [9] C.S. Carvalho and I. Tereno, CMB lensing reconstruction with point source masks, Phys. Rev. D 84, 063001 (2011) (arXiv:1103.2305 [astro-ph])
 - [10] U. Seljak, Gravitational Lensing Effect on Cosmic Microwave Background Anisotropies: A Power Spectrum Approach, Astrophys. J. 463, p. 1 (1996) (arXiv:astro-ph/9505109)
 - [11] M. Bucher, C.S. Carvalho, K. Moodley and M. Remazeilles, CMB lensing reconstruction in real space, Phys. Rev. D 85, 043016 (2012) (arXiv:1004.3285 [astro-ph])
 - [12] We acknowledge D.N. Spergel for suggesting these maps.
 - [13] A. Lewis, A. Challinor and A. Lasenby, Efficient Computation of Cosmic Microwave Background Anisotropies in Closed Friedmann-Robertson-Walker Models, Astrophys. J. 538, p. 473 (2000) (arXiv:astro-ph/9911177)

- [14] K.M. Górski et al, HEALPix: A Framework for High-Resolution Discretization and Fast Analysis of Data Distributed on the Sphere, *Astrophys. J.* 622, p759 (2005) (arXiv:astro-ph/0409513)
- [15] M. Kesden, A. Cooray and M. Kamionkowski, Lensing reconstruction with CMB temperature and polarization, *Phys. Rev. D* 67, 123507 (2003) (arXiv:astro-ph/0302536)
- [16] K.M. Smith, O. Zahn and O. Doré, Detection of Gravitational Lensing in the Cosmic Microwave Background, *Phys. Rev. D* 76, 043510 (2007) (arXiv:0705.3980 [astro-ph])
- [17] S. Pires, S. Plaszczynski and A. Lavabre, Towards a fast, model-independent Cosmic Microwave Background bispectrum estimator, (arXiv:1106.1039 [astro-ph.CO])
- [18] Planck HFI Core Team, Planck early results. VI. The High Frequency Instrument data processing, *A&A*, 536 (2011) (arXiv:1101.2048 [astro-ph.CO]).